arXiv:1207.5892v1 [cond-mat.quant-gas] 25 Jul 2012

Zero-energy states in rotating trapped Bose-Einstein condensates

Tapio Simula
School of Physics, Monash University, Victoria 3800, Australia

We solve the quasiparticle excitation spectrum of a rotating three-dimensional Bose-Einstein condensate whose ground state is pierced by seven singly quantized vortex filaments. We find the elementary excitation spectrum of this system to harbour a quasiparticle with zero excitation energy. This quasiparticle is degenerate with the rotating ground state and may be described as a massless Nambu–Goldstone mode or a Majorana boson. Macroscopic population in such a state corresponds to a Bose–Einstein condensate of Tkachenko vortex wave quanta.

PACS numbers: 03.75.Lm,14.80.Va,67.85.De

The significance of zero-energy excitations in quantum field theories of both elementary and quasiparticles has substantially increased since Majorana predicted the appearance of a mysterious massless particle, which acts as its own anti-particle [1]. Neutrinos have been suggested as candidates for Majorana fermions and more recently, non-elementary Majorana quasiparticles have sparked major interest due to their non-Abelian exchange statistics, which could be harnessed for fault tolerant topological quantum computing [2–4]. Majorana zero modes are also predicted to become trapped inside vortex cores in chiral p-wave paired superconductors and protected by coupling to edge modes [4–6]. Signatures of Majorana modes in superconductor-semiconductor nanowires have recently been observed [7].

Zero-energy excitations have been central to discussions of field theories with spontaneously broken continuous symmetries [8–12]. These massless modes are called Nambu–Goldstone bosons, the photon being a famous example. In non-relativistic condensed matter systems, phase transitions involving spontaneous symmetry breaking, such as the Bose–Einstein condensation phase transition emerge with associated Nambu–Goldstone bosons. In superconductors such zero mode becomes gapped due to the Coulomb interaction [13–15]. Furthermore, the mass of elementary particles is thought to emerge due to shifts in the energies of zero modes [15–20]. Also in systems where continuous symmetries are broken explicitly rather than spontaneously, the Nambu–Goldstone bosons typically acquire small mass correction and are called pseudo-Nambu–Goldstone bosons [21].

Here we study a quantum degenerate Bose gas, which is exposed to an external vector potential that causes the atoms to rotate at an angular velocity Ω around the chosen z-axis of the three-dimensional harmonic potential. By applying the microscopic Bogoliubov–deGennes field theory, we show that this symmetry broken system is capable of hosting a zero mode whose physical appearance is that of a Tkachenko vortex wave. Such an excitation mode is degenerate with the rotating ground state and emerges as a consequence of the broken continuous rotational symmetry in the vortex lattice ground state. Since there is no energy cost associated with populating this excitation mode, we expect these quasiparticles to become macroscopically occupied forming a Bose-Einstein condensate of Tkachenko wave quanta.

By using the method of second quantization, Bogoliubov provided an explanation of the phenomenon of superfluidity in terms of a gas of quasiparticles [22]. In this picture, the quasiparticle creation operator $\hat{\psi}_q$ and annihilation operator $\hat{\psi}_q^\dagger$ obey the equations of motion

$$\begin{aligned} i\hbar\partial_t\hat{\psi}_q &= E_q\hat{\psi}_q \\ i\hbar\partial_t\hat{\psi}_q^\dagger &= -E_q\hat{\psi}_q^\dagger. \end{aligned} \tag{1}$$

Quasiparticles with zero energy $E_q = 0$ satisfy the relation

$$\hat{\psi}_q = \hat{\psi}_q^\dagger, \tag{2}$$

which is the defining property of a Majorana operator—a (quasi) particle creation operator is equivalent to its annihilation operator. The peculiar character of Majorana quasiparticles is not restricted to systems composed of fermionic particles. The Majorana zero mode itself is neither a boson or a fermion as it satisfies neither commutation or anticommutation relations and is therefore described as an anyon [23].

The quasiparticle energies E_q can be explicitly calculated for a given physical system by solving the Bogoliubov–deGennes equations [22, 24, 25]

$$\begin{aligned} \mathcal{L}(\mathbf{r})u_q(\mathbf{r}) - \mathcal{M}(\mathbf{r})v_q(\mathbf{r}) &= (E_q+\mu)u_q(\mathbf{r}) \\ \mathcal{L}^*(\mathbf{r})v_q(\mathbf{r}) - e^{i\theta}\mathcal{M}^*(\mathbf{r})u_q(\mathbf{r}) &= -(E_q-\mu)v_q(\mathbf{r}) \end{aligned} \tag{3}$$

where $\mathcal{L}(\mathbf{r})$ and $\mathcal{M}(\mathbf{r})$ are model dependent operators, $u_q(\mathbf{r})$ and $v_q(\mathbf{r})$ are the quasiparticle amplitudes, and μ is the chemical potential. For the boson system studied here $\theta = 0$, whereas for fermion systems the anticommutation relations yield $\theta = \pi$. For systems whose finite-energy excitations are fermionic the excitation energies E_q are measured with respect to the Fermi energy $E_{\rm F}$, whereas for boson systems they are measured with respect to the chemical potential μ of the condensate.

Considering a Bose–Einstein condensed system, a quadratic approximation to the full many-body Hamiltonian, expressed in terms of the atom field operator $\hat{\Psi}(\mathbf{r})$,



FIG. 1. Nambu–Goldstone boson $\phi(\mathbf{r})$ of a condensed Bose gas rotating at an angular frequency $\Omega = 0.58\omega_\perp$. (a) Isosurface of the condensate density $|\phi(\mathbf{r})|^2$ with seven quantized vortices piercing the condensate forming a sixfold symmetric triangular array, viewed along the rotation axis. (b) The corresponding phase map $S(\mathbf{r}) = \arg(\phi(\mathbf{r}))$ shows the seven phase singularities at the vortex cores, marked with filled circles.

can be diagonalized by the canonical Bogoliubov transformation to a quasiparticle basis [22]

$$\hat{\Psi}(\mathbf{r}) - \phi(\mathbf{r}) = \sum_q u_q(\mathbf{r})\hat{\psi}_q - v_q^*(\mathbf{r})\hat{\psi}_q^\dagger \qquad (4)$$

where $\phi(\mathbf{r})$ is the macroscopic wavefunction describing the atoms in a Bose–Einstein condensate. The quasiparticle creation and annihilation operators $\hat{\psi}_q$ and $\hat{\psi}_q^\dagger$ preserve the canonical boson commutation relations $[\hat{\psi}_q, \hat{\psi}_p^\dagger] = \delta_{qp}$, $[\hat{\psi}_q, \hat{\psi}_p] = [\hat{\psi}_q^\dagger, \hat{\psi}_p^\dagger] = 0$ and the finite energy quasiparticle amplitudes satisfy the orthonormalization condition [24] $\int (u_q^*(\mathbf{r})u_p(\mathbf{r}) - v_q^*(\mathbf{r})v_p(\mathbf{r}))d^3\mathbf{r} = \delta_{qp}$ and $\int (u_q(\mathbf{r})v_p(\mathbf{r}) - u_p(\mathbf{r})v_q(\mathbf{r}))d^3\mathbf{r} = 0$.

We consider a Bose–Einstein condensate of $N = 3\times 10^4$ atoms at zero temperature limit. The ^{87}Rb atoms of mass m are confined in a three-dimensional harmonic oscillator potential $V(\mathbf{r}) = m(\omega_\perp^2 r_\perp^2 + \omega_z^2 z^2)/2$, which rotates at an angular frequency $\Omega = 0.58\omega_\perp$ unless otherwise stated. This rotation frequency stabilizes the seven vortex array as the ground state of this system. The interaction coupling constant $g = 4\pi\hbar^2 a/m$, where a is the s-wave scattering length of the atoms and the oscillator length $a_0 = \sqrt{\hbar/m\omega_\perp}$. We have chosen $\omega_\perp = \omega_z/\sqrt{8} = 2\pi \times 100\,\mathrm{Hz}$. For this system, the operators in Eq. (3) are $\mathcal{M}(\mathbf{r}) = g\phi^2(\mathbf{r})$ and $\mathcal{L}(\mathbf{r}) = -\frac{\hbar^2\nabla^2}{2m} + V(\mathbf{r}) + 2g|\phi(\mathbf{r})|^2 - \Omega L_z$, where L_z denotes the projection of the orbital angular momentum operator onto the axis of rotation. The condensate order parameter $\phi(\mathbf{r})$ satisfies the Gross–Pitaevskii equation [26, 27]

$$\left(\mathcal{L}(\mathbf{r}) - g|\phi(\mathbf{r})|^2 - \mu\right)\phi(\mathbf{r}) = 0. \qquad (5)$$

To draw an analogy with charged superconductors, the operator $\mathcal{L}(\mathbf{r})$ may alternatively be expressed in terms of a vector potential $\mathbf{A}_\Omega = m^2\Omega(-y\mathbf{e}_x + x\mathbf{e}_y)/h$ as $\mathcal{L}(\mathbf{r}) = [(i\hbar\nabla - \kappa\mathbf{A}_\Omega)^2 + m^2(\omega_\perp^2 - \Omega^2)r_\perp^2 + m^2\omega_z^2 z^2 + 4mg|\phi(\mathbf{r})|^2]/2m$, where $\kappa = h/m$ and h is Planck's constant.



FIG. 2. Majorana boson $\psi(\mathbf{r})$ of a condensed Bose gas rotating at an angular frequency $\Omega = 0.58\omega_\perp$. (a) Isosurface of the density $|\psi(\mathbf{r})|^2$ viewed along the rotation axis. (b) Relative phase map $M(\mathbf{r}) = \arg(\phi(\mathbf{r})) - \arg(\psi(\mathbf{r}))$ shows 18 phase singularities. The six outermost of these are ghost vortices of $\psi(\mathbf{r})$ and reside at the edge of the condensate. The remaining 12 phase defects correspond to 6 vortices in the condensate ground state $\phi(\mathbf{r})$ and 6 antivortices in the state $\psi(\mathbf{r})$. Both $\phi(\mathbf{r})$ and $\psi(\mathbf{r})$ have a vortex in the center of the trap. These vortices are of opposite circulation and cancel each other in the relative phase map.

The coupled equations (3) can be shown to possess a particle-hole symmetry such that given a solution $u_q(\mathbf{r}), v_q(\mathbf{r}), E_q$ a second solution $u_q'(\mathbf{r}), v_q'(\mathbf{r}), E_q'$ always exists and is obtained by a simple transformation $u_q'(\mathbf{r}) = v_q^*(\mathbf{r})$, $v_q'(\mathbf{r}) = u_q^*(\mathbf{r})$ and $E_q' = -E_q^*$. For fermion systems both hole and particle type excitations are treated on an equal footing whereas for bosonic systems half of all excitations have a negative norm and in order to avoid describing them as fermionic excitations, they are typically discarded. In addition, for a Bose–Einstein condensed system, there exists a trivial solution, $E_1 = 0$ to Eq. (3), which is obtained by setting $u_1(\mathbf{r}) = v_1^*(\mathbf{r}) = \phi(\mathbf{r})$. This is the Nambu–Goldstone boson corresponding to the spontaneous $U(1)$ gauge symmetry breaking due to the formation of the Bose–Einstein condensate. Such a zero mode does not obey the bosonic quasiparticle normalization condition and instead it satisfies the relation $\int u_1^*(\mathbf{r})u_1(\mathbf{r})d^3\mathbf{r} = \int v_1^*(\mathbf{r})v_1(\mathbf{r})d^3\mathbf{r}$. In addition to the trivial solution, other zero modes degenerate with the ground state and which satisfy $\mathcal{L}(\mathbf{r})u_q(\mathbf{r}) = \mathcal{M}(\mathbf{r})u_q^*(\mathbf{r})$, may exist in the spectrum.

We have calculated the quasiparticle excitation spectra by numerically solving Eq. (3) using a parallelized Arnoldi iteration method. The three dimensional operators are discretized using a finite-element discrete variable representation [28] to yield a sparse matrix representation for the Bogoliubov–de Gennes operator. Before solving the Bogoliubov–de Gennes problem, the condensate ground state $\phi(\mathbf{r})$ and the corresponding chemical potential μ are obtained by solving the Gross–Pitaevskii equation (5) using an over-relaxation method. The self-consistency of the calculated spectrum is guaranteed by the fact that we obtain the $E_1 = 0$ Nambu–Goldstone solution with a numerical precision $|E_1| < 10^{-9}\hbar\omega_\perp$.

Figure 1(a) shows a condensate density $|\phi(\mathbf{r})|^2$ isosurface and (b) the corresponding phase map $S(\mathbf{r})$ for the calculated ground state wavefunction $\phi(\mathbf{r}) = |\phi(\mathbf{r})| \exp(iS(\mathbf{r}))$. The chemical potential of this condensate is $\mu = \langle\phi|\mathcal{L}(\mathbf{r}) - g|\phi(\mathbf{r})|^2|\phi\rangle = 12.0\,\hbar\omega_\perp$ and the orbital angular momentum $L = \langle\phi| - i\hbar(x\partial_y - y\partial_x)|\phi\rangle = 4.2\,\hbar N$. The condensate is pierced by 7 singly quantized vortex filaments. The circulation $\oint v_s(\mathbf{r}) \cdot dl = \kappa\ell$ of the superfluid velocity $v_s(\mathbf{r}) = \hbar/m\nabla S(\mathbf{r})$ around the vortex cores is quantized in integer ℓ multiples of $\kappa = h/m$. In the ground state of the rotating potential, these vortices are arranged in a triangular lattice structure with sixfold discrete rotational symmetry. The phase singularities located inside the vortex cores are marked with filled circles in Fig. 1(b).

Figure 3 shows the Bogoliubov excitation energies calculated for the rotating condensate ground state shown in Fig. 1. Energies of the lowest collective quasiparticle modes are shown as a function of an integer index q, which labels the quasiparticles in increasing order of their energy. The most prominent feature of this excitation spectrum is the presence of two zero energy modes $|E_{1,2}/\hbar\omega_\perp| < 10^{-9}$ labelled by $q = 1$ and $q = 2$. We note that all shown states are duplicated in the numerically calculated spectrum due to the aforementioned particle-hole symmetry of the Bogoliubov–de Gennes equations and thus we find 4 zero modes, only half of which represent distinct quasiparticles and which are shown in Fig. 3. The first one of these modes $q = 1$ is the Nambu–Goldstone boson $\phi(\mathbf{r})$ always present in these Bose–Einstein condensed systems irrespective of the value of the vector potential $\mathbf{A}_\Omega$. It corresponds to the Bose–Einstein condensate of atoms and is equivalent to the state shown in Fig. 1. The presence of this mode reflects the spontaneously broken $U(1)$ gauge symmetry of the ground state as the phase coherent Bose–Einstein condensate is formed.

The observation of the second zero energy mode $q = 2$ corresponds to a massless Nambu–Goldstone mode $\psi(\mathbf{r})$ or a Majorana boson. In the condensate ground state the continuous rotation symmetry SO(2) has also been broken and only the reduced discrete sixfold rotational symmetry of the vortex lattice remains. The source of this symmetry breaking can be traced back to the vector potential $\mathbf{A}_\Omega$ in the Hamiltonian. Therefore one might *a priori* expect to find a zero energy boson mode in the excitation spectrum due to Goldstone's theorem. Instead, the zero mode has been observed to acquire a small mass correction [29]. Furthermore, this zero mode is different from the high angular momentum surface mode, which approaches zero energy when new vortices are nucleated in the system [30, 31].

The physical character of the quasiparticle state $\psi(\mathbf{r})$ is that of a Tkachenko vortex wave [32]. It comprises of elliptically polarized collective motion of the vortices around their equilibrium locations [32–40]. This mode has transverse and longitudinal components corresponding to azimuthal and radial motion of the vortices. The Tkachenko modes of vortex lattices in harmonically trapped Bose–Einstein condensates have been calculated analytically within a vortex continuum approximation and the mode frequencies have been found to approach zero in the centrifugal limit $\Omega \to \omega_\perp$ as the system approaches the quantum-Hall regime [40]. This is in agreement with the experimental observations for large vortex lattices [29]. However, the Tkachenko mode



FIG. 3. Quasiparticle excitation energies as a function of an integer index q which labels the modes in the order of increasing energy. The two zero modes are singled out in the insets. The energy of both of these excitations is $|E_{1,2}/\hbar\omega_\perp| < 10^{-9}$. The center-of-mass Kohn mode is marked with a triangle. The 7 modes whose energies lie between the condensate $q = 1$ and the Kohn mode $q = 8$ are the vortex displacement modes corresponding to the 7 vortex degrees of freedom.



FIG. 4. Quasiparticle probability density of the zero energy Majorana boson mode $\psi(\mathbf{r})$ in the $z = 0$ plane. The six peaks are localized in the vortex cores of the ground state $\phi(\mathbf{r})$. At the origin, there is a phase singularity in both $\phi(\mathbf{r})$ and $\psi(\mathbf{r})$ where the probability density vanishes.

has not been predicted to appear at zero energy for systems with finite $\Omega < \omega_\perp$. Moreover, previous numerical calculations for purely two-dimensional systems [41–44] or three-dimensional prolate systems [45] have not predicted a zero mode other than the condensate Nambu–Goldstone boson in these systems. We have performed calculations for $\Omega/\omega_\perp$ in the range 0-0.8 for the present three-dimensional oblate system and have found a zero energy vortex wave only for the case of seven vortices. In this context, it is amusing that seven classical vortices in a ring arrangement is a marginally stable configuration having a normal mode with precisely zero frequency [46, 47].

We may understand the finite energy of a typical Tkachenko mode $\psi(\mathbf{r})$ in terms of a coupling to the condensate mean-field $\phi(\mathbf{r})$. If $\psi(\mathbf{r})$ absorbs a Nambu–Goldstone boson from $\phi(\mathbf{r})$ it generically changes the orbital angular momentum of the condensate, which leads to a changing vortex density and is tantamount to radial vortex motion. The restoring force provided by the confinement results in the sinusoidal oscillation of $\psi(\mathbf{r})$, which corresponds to a finite excitation energy. When the angular momenta of $\phi(\mathbf{r})$ and $\psi(\mathbf{r})$ are equal, quasiparticle exchange between them may occur without resulting in longitudinal motion of vortices. The finite frequency of a mode which emerges due to a spontaneous symmetry breaking is typically attributed to the Anderson–Higgs mechanism, recently observed in neutral superfluids [48–50]. Here we observe an opposite effect—the excitation frequency of a Tkachenko mode which due to the broken rotation symmetry would be expected to be zero but is nevertheless observed to be finite under normal circumstances, becomes vanishingly small.

The Kohn mode or the center-of-mass dipole mode with frequency $\omega_\perp - \Omega$ has been marked in Fig. 3 by a triangle. The states with $q = 2 - 8$ are the lowest excitation modes in each of the seven Kelvin–Tkachenko vortex wave branches present in this system due to the seven vortex degrees of freedom [45]. The two insets in Fig. 3 show density isosurface plots of the two zero energy states $\phi(\mathbf{r})$ and $\psi(\mathbf{r})$ with $q = 1$ and $q = 2$, respectively. Fig. 2(a) shows a density isosurface plot $|\psi(\mathbf{r})|^2 = |u_2(\mathbf{r})|^2 = |v_2(\mathbf{r})|^2$ viewed along the rotation axis. Fig. 2(b) shows the phase map $M(\mathbf{r}) = \arg(\phi(\mathbf{r})) - \arg(\psi(\mathbf{r}))$. The white markers denote the locations of vortex cores in the Majorana boson state $\psi(\mathbf{r})$ and the black markers correspond to the vortex phase singularities in the condensate ground state. Both $\phi(\mathbf{r})$ and $\psi(\mathbf{r})$ have a vortex phase singularity at the origin, which cancel each other in the relative phase map shown in Fig. 2(b). Finally, figure 4 shows the probability density of the zero energy Majorana boson in the $z = 0$ plane. It can be viewed as a bound state of the effective potential created by the combination of the harmonic trap and the condensate mean-field. There are six prominent peaks which are localized inside the vortex cores of the condensate. There are also seven vortex phase singularities in $\psi(\mathbf{r})$. Six of these are located in between the peaks observed in Fig. 4 and the seventh is located on the z-axis in the center of the figure.

In conclusion, we have found a zero energy quasiparticle state in the elementary excitation spectrum of a rotating Bose–Einstein condensate in a harmonic trap hosting seven singly quantized vortices. The physical character of this zero mode is that of a Tkachenko vortex wave. At zero frequency there is no energy cost associated with populating this excitation mode, suggesting the presence of a Bose–Einstein condensate of quanta of Tkachenko waves. The ground state of this system is doubly degenerate implying a finite entropy of the zero temperature ground state. For $\Omega = \omega_\perp$ the quantum Hall effect is predicted to appear in rotating quantum gases with a lowest Landau level and a massively degenerate ground state. Here we observe ground state degeneracy already at finite $\omega_\perp - \Omega$. There remains an important question regarding the robustness of this zero energy Majorana boson mode with respect to quantum and thermal fluctuations and perturbations in external trapping potential.

In the Feynman interpretation of antiparticles [51, 52], a Majorana zero mode for which a particle and its antiparticle are equivalent could be described in terms of simultaneous forward and backward time propagation of an excitation. Applying this reasoning to the quasiparticle zero mode in our system and noting that time-reversal changes the sense of rotation, we may view the zero energy Tkachenko mode as a quasiparticle which comprises of simultaneous left and right handed rotation.

The zero energy Tkachenko mode has the exceptional character of corresponding to purely transverse motion of the vortices with zero amplitude in the radial component. In contrast, the lowest Tkachenko mode generically has a finite excitation frequency consistent with the longitudinal mode component which emerges due to its coupling to the condensed boson field. This can be compared with the Anderson–Higgs mechanism in superconductors, which is also thought to be responsible for the mass generation of elementary particles [15–20].

I am grateful to Michael Morgan for valuable discussions.

---

[1] E. Majorana, Nuovo Cimento **5**, 171 (1937).

[2] A.Yu. Kitaev, Ann. Phys. **303**, 2 (2003).

[3] A. Y. Kitaev, Phys.-Usp. **44**, 131 (2001).

[4] C. Nayak, S. H. Simon, A. Stern, M. Freedman, and S. Das Sarma, Rev. Mod. Phys. **80**, 1083 (2008).

[5] N. Read and D. Green, Phys. Rev. B **61**, 10267 (2000).

[6] G. E. Volovik, Pisma Zh. Eksp.Teor. Fiz. **70** 601 (1999) ; [JETP Lett. **70** 609 (1999)].

[7] V. Mourik, *et al.* Science **336**, 1003 (2012).

[8] Y. Nambu, Phys. Rev. **117**, 648 (1960).

[9] Y. Nambu and G. Jona-Lasinio Phys. Rev. **122**, 345 (1961).
[10] J. Goldstone, Nuovo Cimento **19**, 154 (1961).
[11] J. Goldstone, A. Salam, and S. Weinberg, Phys. Rev. **127**, 965 (1962).
[12] H. Watanabe and H. Murayama, Phys. Rev. Lett. **108**, 251602 (2012).
[13] P. W. Anderson, Phys. Rev. **110**, 827 (1958).
[14] P. W. Anderson, Phys. Rev. **112**, 1900 (1958).
[15] P. W. Anderson, Phys. Rev. **130**, 439 (1963).
[16] E. C. G. Stueckelberg, Helv. Phys. Acta **11**, 299 (1938).
[17] J. Schwinger, Phys. Rev. **125**, 397 (1962).
[18] P. W. Higgs, Phys. Rev. Lett. **13**, 508 (1964).
[19] F. Englert and R. Brout, Phys. Rev. Lett. **13**, 321 (1964).
[20] G. S. Guralnik, C. R. Hagen, and T. W. B. Kibble, Phys. Rev. Lett. **13**, 585 (1964).
[21] S. Weinberg, Phys. Rev. Lett. **29**, 1698 (1972).
[22] N. N. Bogoliubov, J. Phys. **11**, 23 (1946).
[23] F. Wilczek, Phys. Rev. Lett, **49**, 957 (1982).
[24] A. L. Fetter, Ann. Phys. **70**, 67 (1972).
[25] P. G. de Gennes, Rev. Mod. Phys. **36**, 225 (1964).
[26] L. P. Pitaevskii, Zh. Eksp. Teor. Fiz. **40**, 646 (1961); [Sov. Phys. JETP **13**, 451(1961)].
[27] E. P. Gross, Nuovo Cimento, **20**, 454 (1961).
[28] B. I. Schneider, L.A. Collins, and S. X. Hu, Phys. Rev. E **73**, 036708 (2006).
[29] I. Coddington, P. Engels, V. Schweikhard, and E. A. Cornell, Phys. Rev. Lett. **91**, 100402 (2003).
[30] M. Ueda and T. Nakajima, Phys. Rev. A **73**, 043603 (2006).
[31] D. Dagnino, N. Barberán, M. Lewenstein, J. Dalibard, Nat. Phys **5**, 431 (2009).
[32] V. K. Tkachenko, Zh. Eksp. Teor. Fiz. **49**, 1875 (1965); [Sov. Phys. JETP **22**, 1282 (1966)]; Zh. Eksp. Teor. Fiz. **50**, 1573 (1966); [Sov. Phys. JETP **23**, 1049 (1966)]; Zh. Eksp. Teor. Fiz. **56**, 1763 (1969); [Sov. Phys. JETP **29**, 245 (1969)].
[33] E. S. Raja Gopal, Ann. Phys. (N. Y.) **29**, 350 (1964).
[34] A. L. Fetter, Phys. Rev. B **11**, 2049 (1975).
[35] M. R. Williams and A. L. Fetter, Phys. Rev. B **16**, 4846 (1977).
[36] E. B. Sonin, Zh. Eksp. Teor. Fiz. **70**, 1970 (1976);[Sov. Phys. JETP **43**, 1027 (1976)].
[37] E. B. Sonin, Rev. Mod. Phys. **59**, 87 (1987).
[38] G. Baym and E. Chandler, J. Low Temp. Phys. **50**, 57 (1983).
[39] S. A. Gifford and G. Baym, Phys. Rev. A **70**, 033602 (2004), [Phys. Rev. A 78, 029904(E) (2008)].
[40] G. Baym, Phys. Rev. Lett. **91**, 110402 (2003).
[41] T. P. Simula, A. A. Penckwitt, and R. J. Ballagh, Phys. Rev. Lett. 92, 060401 (2004).
[42] T. Mizushima, Y. Kawaguchi, K. Machida, T. Ohmi, T. Isoshima, and M. M. Salomaa, Phys. Rev. Lett. **92**, 060407 (2004).
[43] S. J. Woo, L. O. Baksmaty, S. Choi, and N. P. Bigelow, Phys. Rev. Lett. **92**, 170402 (2004).
[44] L. O. Baksmaty, S. J. Woo, S. Choi, and N. P. Bigelow, Phys. Rev. Lett. **92**, 160405 (2004).
[45] T. P. Simula and K. Machida, Phys. Rev. A **82**, 063627 (2010).
[46] J. J. Thomson, *Treatise on the motion of vortex rings*, (Macmillan, London 1883).
[47] T. H. Havelock, Phil. Mag. S. 7. **11**, 617 (1931).
[48] U. Bissbort, S. Götze, Y. Li, J. Heinze, J. S. Krauser, M. Weinberg, C. Becker, K. Sengstock, and W. Hofstetter, Phys. Rev. Lett. **106**, 205303 (2011).
[49] L. Pollet and N. Prokofev, Phys. Rev. Lett. 109, 010401 (2012).
[50] M. Endres, T. Fukuhara, D. Pekker, M. Cheneau, P. Schauß, C. Gross, E. Demler, S. Kuhr, I. Bloch, arXiv:1204.5183.
[51] E. C. G. Stueckelberg, Helv. Phys. Acta, **14**, 51 (1941).
[52] R. P. Feynman, Phys. Rev. **76**, 749 (1949).